

Darboux Problem for the Third-Order Bianchi Equation

A. N. Mironov*

Elabuga Institute, Kazan Volga Region Federal University, Kazan, Russia

Received September 30, 2016

Abstract—The existence and uniqueness of the solution of the Darboux problem are proved. The solution of the Darboux problem is constructed in terms of a function similar to the Riemann–Hadamard function.

DOI: 10.1134/S0001434617070069

Keywords: *Darboux problem, Riemann–Hadamard function, Bianchi equation.*

The Darboux problem for a second-order hyperbolic equation with two independent variables has been considered by many authors. For example, let us mention [1, pp. 228–233], [2]–[7].

In the present paper, we study the existence and uniqueness of the solution of the Darboux problem and construct the corresponding Riemann–Hadamard function for the third-order Bianchi equation studied, for example, in [8]–[13]. Darboux-type problems of the Bianchi equation were considered in some particular cases in [9], [11].

1. The equation

$$L(u) \equiv u_{xyz} + a_{110}(x, y, z)u_{xy} + a_{101}(x, y, z)u_{xz} + a_{011}(x, y, z)u_{yz} + a_{100}(x, y, z)u_x + a_{010}(x, y, z)u_y + a_{001}(x, y, z)u_z + a_{000}(x, y, z)u = f(x, y, z) \quad (1)$$

has been called the *third-order Bianchi equation* by several authors [12, p. 6], [13], [14].

We define the function class $C^{(k,l,m)}$ as follows: a function u belongs to the class $C^{(k_1,k_2,k_3)}$ if its continuous derivatives $\partial^{r_1+r_2+r_3}u/\partial x^{r_1}\partial y^{r_2}\partial z^{r_3}$, $r_i = 0, \dots, k_i$, exist. A solution of class $C^{(1,1,1)}$ is said to be regular. Let D be the domain bounded by the planes $x = 0$ (AA_1O_1O), $y = 0$ (AOB), $y = y_0 > 0$ ($A_1O_1B_1$), $z = x$ (OBB_1O_1), $z = z_0 > 0$ (see the figure). We assume that the coefficients of Eq. (1) satisfy the smoothness conditions $a_{ijk} \in C^{(i,j,k)}(\overline{D})$. We let X , Y , and T denote the faces of D for $x = 0$, $y = 0$, and $z = x$, respectively.

Darboux Problem. *In the domain D , to determine the regular solution of Eq. (1) satisfying the boundary conditions*

$$\begin{aligned} u|_{\overline{X}} &= \varphi_1(y, z), & u|_{\overline{Y}} &= \varphi_2(x, z), & u|_{\overline{T}} &= \psi(x, y), \\ \varphi_1(y, 0) &= \psi(0, y), & \varphi_2(x, x) &= \psi(x, 0), & \varphi_1(0, z) &= \varphi_2(0, z), \\ \varphi_1 &\in C^{(1,1)}(\overline{X}), & \varphi_2 &\in C^{(1,1)}(\overline{Y}), & \psi &\in C^{(1,1)}(\overline{T}). \end{aligned} \quad (2)$$

Let us prove the existence and uniqueness of the solution of the Darboux problem. For this, we use the well-known formula for the solution of the Goursat problem [8], [12, p. 28] as a representation of an arbitrary regular solution of Eq. (1). From this formula we derive the Volterra integral equation of the second kind which we use to pose the Goursat condition $u(x, y, z_0)$ on the plane $z = z_0$ and note that the existence and uniqueness of the solution of this problem imply the existence and uniqueness of the solution of the Darboux problem (1), (2).

*E-mail: lbmironova@yandex.ru